

Ocular Signs in Fabry Disease Case report

Khamaily M*, Salem JB, Tarib I, Sidi D, Bajjouk S, Zerrouk R, Mouzari Y, Asri1 FE, Reda K and Oubaaz A

Department of Ophthalmology, Faculty of Medicine and Pharmacy, Mohammed V University Rabat, Morocco

Volume 4 Issue 7- 2020

Received Date: 14 May 2020

Accepted Date: 04 June 2020

Published Date: 09 June 2020

2. Key words

α galactosidase A; cornea verticillata;
Hereditary; Far by disease

1. Case Report

1.1. Abstract: Fabry disease is a rare, hereditary disease characterized by a deficiency of an enzyme, α galactosidase A (α gal A), responsible for progressive damage to many organs, leading to various symptoms, Ocular damage, particularly to the cornea, is sometimes a precious element helping the positive diagnosis of the disease. We report the case of a 40-year-old patient diagnosed with Fabry disease, with bilateral conjunctival vascular tortuosity's, a "cornea verticillata and a peripheral cortical cataract.

Better knowledge of ophthalmological signs, allows better screening and can participate in the evaluation of the effectiveness of substitute therapy.

3. Introduction

Fabry disease is a rare, hereditary disease, the transmission of which is linked to the X chromosome. It was first described in 1898 by Johannes Fabry and William Anderson [1].

It is characterized by a deficiency of an enzyme, α galactosidase A (α gal A), contained in lysosomes. This deficit causes the accumulation within the lysosomes of a glycosphingolipid, globotriaosylceramide (Gb3), responsible for progressive damage to many organs, leading to various symptoms: ocular, digestive, cutaneous, cardiac, renal, pulmonary but also involvement of the central and peripheral nervous system [2]. The disease develops gradually and the diagnosis is often made late.

Ocular damage, particularly in the cornea, is sometimes a precious element helping the positive diagnosis of the disease.

4. Materials and Methods

We report a case of a 40-year-old patient, diagnosed with hypertrophic heart disease, and Fabry disease since 2012, and who consults in ophthalmology as part of his general check.

5. Results

The patient underwent a complete ophthalmological examination which revealed better visual acuity corrected to 16/20 OSD, a normal shirmer test, in the slit lamp: bilateral conjunctival vascular tortuosity's (Figure 1), with a "cornea verticillata and a peripheral cortical cataract (Figure 2), normal Intra Ocular Pressure and fundus with no vascular dilatation.




Figure 1: Tortuosities of the conjunctival vessels in the two eyes (arrows)




Figure 2: Cornea verticillata (stars), Peripheral cortical cataract (arrow)

6. Discussion

The most distinctive and common clinical feature of FD (90%) is whorl-shaped bilateral keratopathy [3], known as cornea verticillata, secondary to Gb-3 deposits at the level of the epithelial basement membrane, and visualized as yellowish brown inclusions emanating radially; the corneal stroma and the endothelium are spared [4].

Confocal microscopies have a great interest in the detection of sub-clinical lesions and can represent a reliable tool for the early diagnosis and the follow-up of the disease [5].

In children, the presence of eye signs, especially the cornea verticillata, is correlated with a severe form of the Fabry disease [6].

*Corresponding Author (s): Mehdi Khamaily, Department of Ophthalmology, Faculty of Medicine and Pharmacy, Mohammed V University Rabat, Morocco, E-mail: mehdi.khamaily@gmail.com

Conjunctival involvement is a tortuosity of the conjunctival vessels secondary to a disturbance of the endothelial architecture by accumulation of Gb-3.

The lens involvement is represented by two types of clouding, the "classic" Fabry cataracts which result from deposits of Gb-3 in the lens epithelium and which are better visible in back-illumination, which appear as cloudiness under -capsular or dendritic away from the visual axis, and cataracts in "Helices" which are in rays oriented radially in the equatorial region or in subcapsular [7].

These cataracts appear in the second decade of life and are seen in up to 70% of men, less often in women [8].

Dry eyes are present in approximately 50% of patients with Fardy's disease, following deposits of Gb-3 in the lymph nodes and directly in the lacrimal gland [9].

Other eye damage described but much less rare, corneal edema, conjunctival chemosis, and chronic uveitis [10].

7. Conclusion

Fabry's disease remains potentially serious due to its systemic damage, and despite the frequency of ocular damage, which is a precious tool for positive diagnosis, the visual prognosis is very rarely involved.

Better knowledge of ophthalmological signs, allows better screening and can participate in the evaluation of the effectiveness of substitute therapy.

References

1. Sher NA, Letson RD, Desnick RJ. The ocular manifestations in Fabry's disease. *Arch Ophthalmol.* 1979; 97(4): 671-6.
2. Pinakin Gunvant Davey. Fabry disease: a survey of visual and ocular symptoms. *Clinical Ophthalmology.* 2014; 8: 1555-1560.
3. Langis Michaud I. Longitudinal study on ocular manifestations in a cohort of patients with Fabry disease. *PLoS ONE.* 2019; 14(6): e0213329.
4. Morier AM, Minter J, Tyszko R, McCann R, Clarke MV, Browning MF. Ocular manifestations of Fabry disease within in a single kindred. *Optometry.* 2010; 81:437Y49.
5. Andrea Leonardi. Cornea verticillata in Fabry disease: a comparative study between slit-lamp examination and in vivo corneal confocal microscopy. *Br J Ophthalmol.* 2020; 104(5): 718-722.
6. Gisela Kalkum, Susanne Pitz, Nesrin Karabul, Michael Beck, Guillem Pintos-Morell, Rossella Parini, et al. Paediatric Fabry disease: prognostic significance of ocular changes for disease severity *BMC Ophthalmology.* 2016; 16: 202.
7. Sodi A, Ioannidis AS, Mehta A, Davey C, Beck M, Pitz S. Ocular

manifestations of Fabry's disease: data from the Fabry Outcome Survey. *Br J Ophthalmol.* 2007; 91(2): 210-4.

8. Orssaud C, Dufier J, Germain D. Ocular manifestations in Fabry disease: a survey of 32 hemizygous male patients. *Ophthalmic Genet.* 2003; 24(3): 129-39.
9. Katherine Rothstein, Jubby M. Gálvez, Ángela M. Gutiérrez, Laura Rico, Eveling Criollo, Alejandra de-la-Torre. Ocular findings in Fabry disease in Colombian patients. *Biomédica.* 2019; 39(3): 434-39.
10. S I Kharlap, M S Kharlap, A R Salikhova, T S Mitichkina, Z V Surnina, N V Miroshnik, S E Avetisov. Clinical and morphological changes in the eye in Fabry's disease. *Oftalma.* 2019; 135(3): 78-89.